arXiv:0709.4449v1 [math-ph] 27 Sep 2007

On soliton structure of higher order (2+1)-dimensional equations of a relaxing medium beneath high-frequency perturbations

Kuetche Kamgang Victor,[1,*] Bouetou Bouetou Thomas,[2,3,†] and Timoleon Crepin Kofane[1,3,‡]

[1] *Department of Physics, Faculty of Science, University of Yaounde I, P.O. Box. 812, Cameroon*
[2] *Ecole Nationale Supérieure Polytechnique, University of Yaounde I, P.O. Box. 8390, Cameroon*
[3] *The Abdus Salam International Centre for Theoretical Physics,*
P.O. Box 586, Strada Costiera, II-34014, Trieste, Italy

(Dated: February 2, 2008)

We investigate the soliton structure of novel (2+1)-dimensional nonlinear partial differential evolution (NLPDE) equations which may govern the behavior of a barothropic relaxing medium beneath high-frequency perturbations. As a result, we may derive some soliton solutions amongst which three typical pattern formations with loop-, cusp- and hump-like shapes.

PACS numbers: 02.30.Ik, 05.45.Yv

In real natural world, there exist very complicated phenomena closely related to nonlinear systems. However, in various cases, the real natural phenomena are too intricate to be described only by virtue of (1+1)-dimensional NLPDE equations. Thus, a great deal of interest has recently been paid to higher dimensional NLPDE equations, especially, the (2+1)-dimensional cases ([1] and references therein). Such higher dimensional systems have been shown to possess soliton solutions which may arise as a balance between nonlinearity and dispersion. One underlying query that may arise, consists of the survival of the soliton properties of solutions to such systems with higher order nonlinearity. In the wake of such query, we consider a barothropic relaxing medium $p = p(\rho, \lambda)$ under high-frequency perturbations $p' \ll p_0$. The quantities p and ρ may stand for pressure and mass density, respectively. The constant λ is an additional parameter. The pressure p_0 is measured at the unperturbed state. For $\lambda = 1$, $p = p_f$, inner interaction processes are frozen and the velocities for fast processes may be defined as $v_f^2 = \frac{dp_f}{d\rho}$. For $\lambda = 0$, $p = p_e$, there is local thermodynamic equilibrium and the velocities for slow processes may be defined as $v_e^2 = \frac{dp_e}{d\rho}$.

A few years ago, Danylenko et al. [2] have proposed dynamical equations of state for multicomponent relaxing media. In the wake of such result, after a suitable expansion of the specific volume $V \equiv \rho^{-1}$ as power series of perturbation p' with accuracy $o(p'^3)$, the following dynamical equation may be written down

$$\tau\partial_t\left(\partial^2_{x_m x^m}p' - \frac{1}{v_f^2}\partial_t^2 p' + \alpha_f\partial_t^2 p'^2 + a_f\partial_t^2 p'^3\right) + \partial^2_{x_m x^m}p' - \frac{1}{v_e^2}\partial_t^2 p' + \alpha_e\partial_t^2 p'^2 + a_e\partial_t^2 p'^3 = 0, \quad (m = 1, \cdots, N), \quad (1)$$

where τ is the relaxation time; constants α_e and α_f represent positive-valued second order expansion cœfficients of V_e and V_f, respectively; constants a_e and a_f stand for positive-valued third order expansion cœfficients of V_e and V_f, respectively; quantity $x^m \equiv (x, y, z, \cdots)$ may stand for position-vector and $\partial^2_{x_m x^m} = \partial^2_{x_1 x^1} + \partial^2_{x_2 x^2} + \partial^2_{x_3 x^3} + \cdots$, $(m = 1, \cdots, N)$. It is noted that any repeated index refers to summation with respect to Einstein's notation. In order to investigate the eq. (1), the multiscale method [3, 4] may be useful. Thus, defining the quantity $\epsilon = \tau\omega$ (ω being the frequency of the processes) chosen to be small (large) parameter, after introducing the following independent variables $T_0 = t\omega$, $T_{-4} = t\omega\epsilon^{-4}$, $X_0^m = x^m\omega$, $X_{-4}^m = x^m\omega\epsilon^{-4}$, $(m = 1, \cdots, N)$ into eq. (1), seven coupled equations may be derived. From this coupled system, there may be two leading equations expressed in terms of T_0 and X_0^m $(m = 1, \cdots, N)$, describing low-frequency perturbations, and the two other expressed in terms of T_{-4} and X_{-4}^m $(m = 1, \cdots, N)$ describing high-frequency perturbations. Thus, focusing our interest only to high frequency perturbations, we may derive the following evolution equation

$$(\partial_{x_n} + \partial_{x_m})(\partial_{x_n} + \partial_{x_m})\left(p' + \alpha_f v_f^2 p'^2 + a_f v_f^2 p'^3\right) - v_f^{-2}\partial_{tt}p' + \beta_f\partial_{x_m}J^m + \gamma_f p' = 0, \quad (n < m), \quad (2)$$

where $J^m = p'$ $(m = 1, \cdots, N)$, and the quantities β_f and γ_f may be expressed as follows

$$\beta_f = \frac{v_f^2 - v_e^2}{\tau v_e^2 v_f}, \quad \gamma_f = \frac{v_f^4 - v_e^4}{2\tau^2 v_e^4 v_f^2}. \quad (3)$$

---

[*] Electronic address: vkuetche@yahoo.fr
[†] Electronic address: tbouetou@yahoo.fr
[‡] Electronic address: tckofane@yahoo.com

Eq. (2) may be obtained in the following way. A dispersion relation for the linearized eq. (2) may be written down in the form $v_f^{-2}\omega^2 = (k_n + k_m)(k_n + k_m) + j\beta_f \sum_{m=1}^{N} k_m - \gamma_f$ with $(n < m = 1, \cdots, N)$. The nonlinear terms may be reconstructed in agreement with the initial equation. We may note here that eq. (2) has dissipative β_f-terms and dispersive γ_f-terms. If $\alpha_f = a_f = \beta_f = 0$, eq. (2) reduces to a typical modified Klein-Gordon equation in (N+1)-dimensional space. We particularly focus our interests to $N = 2$ case. Then, it comes

$$(\partial_x + \partial_y)^2 p' - v_f^{-2}\partial_{tt}^2 p' + \alpha_f v_f^2 (\partial_x + \partial_y)^2 p'^2 + a_f v_f^2 (\partial_x + \partial_y)^2 p'^3 + \beta_f (\partial_x + \partial_y) p' + \gamma_f p' = 0. \tag{4}$$

If $a_f = 0$ and y-terms removed, eq. (4) may reduce to the (1+1)-dimensional Vakhnenko (V) equation [5, 6, 7, 8, 9]. We need the following accuracy

$$(\partial_x + \partial_y)^2 - v_f^{-2}\partial_{tt}^2 \approx 2(\partial_x + \partial_y)\left(\partial_x + \partial_y + v_f^{-1}\partial_t\right), \tag{5}$$

to further investigate the eq. (4). Thus, it may be interesting to consider two important cases: $\alpha_f = 0$ and $\alpha_f \neq 0$.

1. Case $\alpha_f = 0$,

   then, eq. (4) may be transformed to

   $$(\partial_{\tilde{x}} + \partial_{\tilde{y}})\left[\partial_{\tilde{t}} - \frac{1}{2}\tilde{u}^2(\partial_{\tilde{x}} + \partial_{\tilde{y}})\right]\tilde{u} + \tilde{\alpha}(\partial_{\tilde{x}} + \partial_{\tilde{y}})\tilde{u} - \tilde{u} = 0, \tag{6}$$

   provided

   $$\tilde{x} = -\sqrt{\frac{\gamma_f}{6}}(x - v_f t), \quad \tilde{y} = -\sqrt{\frac{\gamma_f}{6}}(y - v_f t), \quad \tilde{t} = v_f\sqrt{\frac{3\gamma_f}{2}}t, \quad \tilde{u} = \alpha_f v_f^2 p', \quad \tilde{\alpha} = \frac{\beta_f}{\sqrt{6\gamma_f}}, \tag{7}$$

   hold. Performing variable transformation, we introduce new independent variables X, T_1 and T_2 as follows

   $$\tilde{x} = T_1 - \frac{1}{2}\int_{-\infty}^{X} U^2 dX' + \tilde{x}_0, \quad \tilde{y} = T_2 - \frac{1}{2}\int_{-\infty}^{X} U^2 dX' + \tilde{y}_0, \quad \tilde{t} = X, \tag{8}$$

   where $\tilde{x}_0$ and $\tilde{y}_0$ stand for arbitrary constants and $\tilde{u}(\tilde{x}, \tilde{y}, \tilde{t}) = U(T_1, T_2, X)$. Eq. (6) may be rewritten as follows

   $$U_{XT_1} + U_{XT_2} + \tilde{\alpha}(U_{T_1} + U_{T_2}) - (1 + \varphi + \phi)U = 0, \tag{9}$$

   where $\varphi = -\int_{-\infty}^{X} UU_{T_1} dX'$ and $\phi = -\int_{-\infty}^{X} UU_{T_2} dX'$. Subscripts with respect to X, T_1 and T_2 may denote partial differentiations. Using the ansates $\varphi = Z_{1T_1} - 1$ and $\phi = Z_{2T_2} - 1$, eq. (6) may bilinearize to

   $$(D_X D_{T_1} + D_X D_{T_2} + \tilde{\alpha}(D_{T_1} + D_{T_2}) - 1)G \cdot F = 0, \quad D_X^2 F \cdot F = \frac{1}{2}G^2, \tag{10}$$

   provided

   $$U = \frac{G}{F}, \quad Z_1 = T_1 - 2\left(\ln(F)\right)_X, \quad Z_2 = T_2 - 2\left(\ln(F)\right)_X, \tag{11}$$

   hold. Notations D_X, D_{T_1} and D_{T_2} denote Hirota operators [10, 11]. Expanding F and G in a suitable formal power series, a one-soliton solution to eq. (9) may be given by

   $$U = 2Ksech(\theta), \quad Z_1 = T_1 - 2K\left[\tanh(\theta) + 1\right], \quad Z_2 = T_2 - 2K\left[\tanh(\theta) + 1\right]. \tag{12}$$

   where $\theta = KX - \omega_1 T_1 - \omega_1 T_1 + \theta_0$, θ_0 being an arbitrary constant. The dispersion relation may be given by

   $$(K + \tilde{\alpha})(\omega_1 + \omega_2) + 1 = 0. \tag{13}$$

   As a result, assuming that $\omega_1 = Kv_1$ and $\omega_2 = Kv_2$ such that $v = v_1 + v_2 < 0$, we may find out three typical pattern formations. Indeed,

   (a) for $\tilde{\alpha} < \sqrt{\frac{1}{2|v|}}$, loop-like pattern may be obtained (see FIG. 1 in the case of $v = -0.24$);

   (b) for $\tilde{\alpha} = \tilde{\alpha}_c = \sqrt{\frac{1}{2|v|}}$, cusp-like pattern may be obtained (see FIG. 2 in the case of $v = -0.24$ where $\tilde{\alpha}_c = 1,4433756729740644112728719512549$);

   (c) finally, for $\tilde{\alpha} > \sqrt{\frac{1}{2|v|}}$, hump-like pattern may be obtained (see FIG. 3 in the case of $v = -0.24$).

These illustrative curves are plotted at initial time $\tilde{t} = 0$. An essential remark on eq. (6) should be noted here. In the wake of the results got from ref. [12], an extended complex-valued form of eq. (6) may be found as follows

$$(\partial_{\tilde{x}} + \partial_{\tilde{y}}) \left[\partial_{\tilde{t}} - \frac{1}{2} \tilde{q}\tilde{q}^{\star}(\partial_{\tilde{x}} + \partial_{\tilde{y}}) \right] \tilde{q} + \tilde{\alpha}(\partial_{\tilde{x}} + \partial_{\tilde{y}})\tilde{q} - \tilde{q} = 0, \tag{14}$$

where $q = q^r + \imath q^{im}$ may stand for a complex-valued observable, symbol $(\star)$ may refer to complex conjugation and $\imath^2 = -1$. Thus, one may derive the following set

$$Q^r_{XT_1} + Q^r_{XT_2} + \tilde{\alpha} \left(Q^r_{T_1} + Q^r_{T_2} \right) - \left(1 + Z_{1T_1} + Z_{2T_2} \right) Q^r = 0, \tag{15a}$$

$$Q^{im}_{XT_1} + Q^{im}_{XT_2} + \tilde{\alpha} \left(Q^{im}_{T_1} + Q^{im}_{T_2} \right) - \left(1 + Z_{1T_1} + Z_{2T_2} \right) Q^{im} = 0, \tag{15b}$$

$$Z_{1XT_1} = - \left(Q^r Q^r_{T_1} + Q^{im} Q^{im}_{T_1} \right), \quad Z_{1XT_2} = - \left(Q^r Q^r_{T_2} + Q^{im} Q^{im}_{T_2} \right), \tag{15c}$$

where $\tilde{q}(\tilde{x}, \tilde{y}, \tilde{t}) = Q(T_1, T_2, X)$ such that independent variables $\tilde{x}$, $\tilde{y}$ and $\tilde{t}$ now stand for

$$\tilde{x} = T_1 - \frac{1}{2} \int_{-\infty}^{X} QQ^{\star} dX' + \tilde{x}_0, \quad \tilde{y} = T_2 - \frac{1}{2} \int_{-\infty}^{X} QQ^{\star} dX' + \tilde{y}_0, \quad \tilde{t} = X. \tag{16}$$

It comes that eq. (15) may be closely related to the coupled dispersionless systems [13, 14] recently investigated by Kakuhata and Konno [15, 16]. Besides, eq. (15) may be also observed as a coupled (2+1)-dimensional version of the complex-form of the Schäfer-Wayne short pulse (SWSP) equation [17] that has been subject to many recent investigations [12, 18, 19, 20, 21]. A one-soliton solution to eq. (15) may be written as follows

$$Q = A \exp(\theta^{im}) sech(\theta^r), \quad Z_1 = T_1 - 2K^r \left[\tanh(\theta^r) + 1 \right], \quad Z_2 = T_2 - 2K^r \left[\tanh(\theta^r) + 1 \right], \tag{17}$$

where $\theta = \theta^r + \imath\theta^{im} = KX - T_1\omega_1 - T_2\omega_2 + \theta_0$, θ_0 being an arbitrary complex-valued constant, $K = K^r + \imath K^{im}$ and $\omega_j = \omega^r_j + \imath\omega^{im}_j$ $(j = 1, 2)$. The amplitude A may be given by $A = 2K^r$. The corresponding dispersion relation may be given by eq. (13). As a result, one may easily find that solutions given by eq. (17) in terms of Q^r and Q^{im}, possess a nonzero angular momentum i.e. the previous patterns depicted above may rotate with an angular frequency $\Omega = \omega^{im}_1 + \omega^{im}_2$. Concretely, if we assume that $\omega^r_1 = v_1 K^r$, $K^{im} = \zeta\omega^{im}_1$, $\omega^r_2 = v_2 K^r$, $K^{im} = \zeta\omega^{im}_2$ and $v = v_1 + v_2 < 0$, K^r and Ω may be expressed as follows

$$K^r = \frac{\tilde{\alpha}}{\zeta|v| - 1}, \quad \Omega = \frac{1}{\sqrt{\zeta}} \sqrt{1 - \frac{\zeta\tilde{\alpha}^2 v^2}{(\zeta v - 1)^2}}, \tag{18}$$

provided $\frac{1}{|v|} < \zeta < \frac{2}{|v|}$. Thus, setting $\alpha_c = \frac{\zeta|v|-1}{\sqrt{2|v|}}$ and $\tilde{\alpha}_s = \frac{\zeta|v|-1}{|v|\sqrt{\zeta}}$, it comes that,

   (a) for $\alpha \geq \tilde{\alpha}_s$, no rotating pattern formation is expected;

   (b) for $\tilde{\alpha}_c < \alpha < \tilde{\alpha}_s$, rotating loop-like pattern formation is derived (see FIG.1 for $v = -0.24$ and $\zeta = 13/3$);

   (c) for $\tilde{\alpha} = \alpha_c$, rotating cusp-like pattern formation is derived (see FIG.2 for $v = -0.24$ and $\zeta = 13/3$ where $\alpha_c = 0,057735026918962576450914878050196$);

   (d) finally, for $\alpha < \tilde{\alpha}_c$, rotating hump-like pattern formation is derived (see FIG.3 for $v = -0.24$ and $\zeta = 13/3$).

2. Case $\alpha_f \neq 0$,

   then, eq. (4) may lead to

$$(\partial_{\tilde{x}} + \partial_{\tilde{y}}) \left[\partial_{\tilde{t}} + \tilde{u}(\partial_{\tilde{x}} + \partial_{\tilde{y}}) + \frac{1}{2} \tilde{u}^2 (\partial_{\tilde{x}} + \partial_{\tilde{y}}) \right] \tilde{u} + \tilde{\alpha}(\partial_{\tilde{x}} + \partial_{\tilde{y}})\tilde{u} + \tilde{u} = 0, \tag{19}$$

provided

$$\tilde{x} = \sqrt{\frac{3a_f\gamma_f}{2\alpha_f^2 v_f^2}}(x - v_f t), \quad \tilde{y} = \sqrt{\frac{3a_f\gamma_f}{2\alpha_f^2 v_f^2}}(y - v_f t), \quad \tilde{t} = \sqrt{\frac{\gamma_f}{6a_f}}\alpha_f v_f^2 t, \quad \tilde{u} = \frac{3a_f}{\alpha_f}p', \quad \tilde{\alpha} = \frac{\beta_f}{\alpha_f v_f}\sqrt{\frac{3a_f}{2\gamma_f}}, \tag{20}$$

hold. We introduce new independent variables X', T_1' and T_2' as follows

$$\tilde{x} = T_1' + \frac{1}{2}\int_{-\infty}^{X'} \left(U^2 + 2U\right) dS + \tilde{x}_0', \quad \tilde{y} = T_2' + \frac{1}{2}\int_{-\infty}^{X'} \left(U^2 + 2U\right) dS + \tilde{y}_0', \quad \tilde{t} = X', \tag{21}$$

where $\tilde{x}_0'$ and $\tilde{y}_0'$ stand for arbitrary constants and $\tilde{u}(\tilde{x}, \tilde{y}, \tilde{t}) = U(T_1', T_2', X')$. Then eq. (19) becomes

$$U_{X'T_1'} + U_{X'T_2'} + \tilde{\alpha}(U_{T_1'} + U_{T_2'}) + (1 + \varphi' + \phi')U = 0, \tag{22}$$

where $\varphi' = \int_{-\infty}^{X'}(U+1)U_{T_1}dS$ and $\phi' = \int_{-\infty}^{X}(U+1)U_{T_2}dS$. Using the ansates $\varphi' = -Z_{1T_1'}+1$ and $\phi' = -Z_{2T_2'}+1$, eq. (19) may be bilinearized as

$$(D_{X'}D_{T_1'} + D_{X'}D_{T_2'} + \tilde{\alpha}(D_{T_1'} + D_{T_2'}) + 1)G \cdot F = 0, \quad D_X^2 F \cdot F = \frac{1}{2}\left(G^2 + 2GF\right), \tag{23}$$

provided

$$U = \frac{G}{F}, \quad Z_1 = T_1' - 2\left(\ln(F)\right)_{X'}, \quad Z_2 = T_2' - 2\left(\ln(F)\right)_{X'}, \tag{24}$$

hold. Thus expanding suitably the functions G and F in power series according to Hirota method [10, 11], one-soliton solutions of kink-like pattern may be expected and discussed by means of the dispersion relation derived from eq. (23).

In summary, we have given a general model of a (2+1)-dimensional NLPDE equation (see eq. (14)) with the soliton structure. This novel (2+1)-dimensional NLPDE equation (eq. (14)) may be observed as a coupled (2+1)-dimensional version of the (1+1)-dimensional complex-SWSP equation [12]. This novel equation with the other one (see eq. (19)) may deserve further scientific interests both from the viewpoint of the investigation of the propagation of high-frequency perturbations and from the viewpoint of the existence of stable wave formations. We may actually believe that these equations may be valuable for studies on soliton theory, geodynamics, plasma physics, hydrodynamics, condensed matter, string theory, nonlinear optics, just to name a few.

---

[1] X. Y. Tang, and S. Y. Lou, J. Math. Phys. **44**, 4000 (2003).
[2] V. A. Danylenko, V. V. Sorokina, and V. A. Vladimirov, J. Phys. A **26**, 7125 (1993).
[3] A. H. Nayfey, *Perturbation methods*, (Wiley, New-York, 1973).
[4] Y. A. Nitropolsky, A. M. Samoilenko, and D. I. Vladimirov, *Systems of evolution equations with periodic and quasiperiodic cœfficients*, (Kluwer Academic, Dordrecht, 1993).
[5] V. O. Vakhnenko, J. Math. Phys. **40**, 2011 (1999).
[6] V. O. Vakhnenko and E. J. Parkes, Nonlinearity **11**, 1457 (1998).
[7] V. O. Vakhnenko, E. J. Parkes, and A. J. Morrison, Chaos Solitons Fractals **17**, 683 (2003).
[8] A. J. Morrison and E. J. Parkes, Chaos Solitons Fractals **16**, 13 (2003).
[9] A. J. Morrison and E. J. Parkes, Glasgow Math. J. **43**, 65 (2001).
[10] R. Hirota, *Solitons*, (Springer, New-York, 1980).
[11] R. Hirota, *Direct methods in soliton theory*, (Springer-Verlag, Berlin, 1988).
[12] V. K. Kuetche, T. B. Bouetou, and T. C. Kofane, J. Phys. Soc. Jpn. **76**, 073001 (2007).
[13] T. Alagesan and K. Porsezian, Chaos Solitons Fractals **8**, 1645 (1997).
[14] V. K. Kuetche, T. B. Bouetou, and T. C. Kofane, J. Phys. A **39**, 12355 (2006).
[15] H. Kakuhata and K. Konno, J. Phys. Soc. Jpn. **68**, 757 (1999).
[16] H. Kakuhata and K. Konno, Theor. Math. Phys. **133**, 1675 (2002).
[17] T. Schäfer and C. E. Wayne, Physica D **196**, 90 (2004).
[18] E. J. Parkes, Chaos Solitons Fractals: in press, (2006).
[19] V. K. Kuetche, T. B. Bouetou, and T. C. Kofane, J. Phys. A **40**, 5585 (2007).
[20] A. Sakovich and S. Sakovich, Glasgow Math. J. **39**, L361 (2006).
[21] V. K. Kuetche, T. B. Bouetou, and T. C. Kofane, J. Phys. Soc. Jpn. **76**, 024004 (2007).